

GiNaCDE: the high-performance F-expansion and First Integral Methods with C++ library for solving Nonlinear Differential Equations

Mithun Bairagi¹

¹ Department of Physics, The University of Burdwan, Golapbag 713104, West Bengal, India

DOI: [10.21105/joss.03885](https://doi.org/10.21105/joss.03885)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Marie E. Rognes](#) ↗

Reviewers:

- [@peanutfun](#)
- [@carstenbauer](#)

Submitted: 12 September 2021

Published: 21 April 2022

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

GiNaCDE is a free and open-source C++ library that solves NLPDEs (Nonlinear Partial Differential Equations) and NLODEs (Nonlinear Ordinary Differential Equations) based on the high-performance F-expansion (Zhou et al., 2003), modified F-expansion (mF-expansion) (Cai et al., 2006) and First Integral Methods (FIM) (Feng, 2002). It unifies and gathers various research on the F-expansion method and FIM (Bhrawy et al., 2013; Cai et al., 2006; Feng, 2002; He, 2013; Huang et al., 2007; Mirzazadeh & Eslami, 2012; Na, 2004; Zhang et al., 2010; Zhou et al., 2003). It implements two versions of the F-expansion method: F-expansion and modified F-expansion methods. It can be used to get exact traveling-wave solutions of a wide variety of NLPDEs arising in different scientific fields by applying any one of the three available methods (F-expansion, mF-expansion, and first integral method). GiNaCDE is built on a pure C++ symbolic library GiNaC (Bauer et al., 2002), and its algebraic manipulations are performed by the classes provided by the GiNaC library. It also has a rich Graphical User Interface (GUI) which makes it more user-friendly.

The library has been designed to solve the NLPDEs, which have the following general form

$$F(\alpha_i, u, u_t, u_{x_1}, u_{x_1} \dots, u_{x_m}, u_{tt}, u_{tx_1}, u_{tx_2}, \dots, u_{tx_m}, u_{x_1x_1}, u_{x_1x_2} \dots, u_{x_1x_m} \dots) = 0, \quad (1)$$

where t, x_1, x_2, \dots, x_m are independent variables, $u = u(t, x_1, x_2, \dots, x_m)$ is a dependent variable, $\alpha_i (i = 1, 2, \dots, n)$ are the parameters. Here F must be a polynomial in u and its derivatives. The primary intention of GiNaCDE is to facilitate the development and validation of new exact analytical solutions of NLPDEs and NLODEs. Using the GiNaCDE library, we have successfully solved various kinds of NLPDEs, including higher nonlinearity terms, higher-derivative terms, and complex NLPDE. Some of them are: Eckhaus equation, seventh-order Sawada-Kotara equations, fifth-order Generalized Korteweg–De Vries (KdV) equation, perturbed nonlinear Schrödinger (NLS) equation with Kerr Law Nonlinearity, and Kudryashov-Sinelshchikov equation.

Statement of need

The NLODEs and NLPDEs play an important role in the theoretical sciences to explain many nonlinear phenomena in various fields of science, such as biology, chemistry, engineering, solid-state physics, plasma physics, optical fibers. The exact (closed-form) traveling-wave solutions of such NLPDEs give much extra information, which helps us to study the result more deeply. The knowledge of closed-form solutions of NLODEs and NLPDEs helps to test the degree of accuracy of numerical solvers and also facilitates stability analysis. In the past few decades, many powerful methods have been presented to seek exact solutions of NLPDEs, such as F-expansion method and first integral method. The F-expansion method was first

proposed by Zhou et al. (Zhou et al., 2003). The first integral method was first introduced by Feng (Feng, 2002) in solving the Compound Burgers-KdV Equation, which is based on the ring theory of commutative algebra. Later, these methods have been further improved in a number of research works. Following some research works (Bhrawy et al., 2013; Cai et al., 2006; Feng, 2002; He, 2013; Huang et al., 2007; Mirzazadeh & Eslami, 2012; Na, 2004; Zhang et al., 2010; Zhou et al., 2003), GiNaCDE intends to gather and unify the possible combinations of the different revised versions of these methods for improvements of solution procedures. In this context, on the basis of the aforementioned research works of these methods, we have presented the high-performance algorithms of F-expansion, modified F-expansion, and first integral methods in the documentation¹ file. The GiNaCDE software uses these algorithms to solve the NLPDEs and NLODEs.

In order to solve the NLPDEs, many computer packages are available. In 1996, Parkes and Duffy (Parkes & Duffy, 1996) had implemented tanh-expansion in their Mathematica package ATFM. Later complete implementation of tanh-expansion has been done by Li and Liu (2002) (Li & Liu, 2002) designing the Maple package RATH. Baldwin et al. (Baldwin et al., 2004) have developed the Mathematica package *PDESpecialSolutions.m* which admits polynomial solutions in tanh, sech, combinations thereof, JacobiSN, JacobiCN. RAEEM (Lin & Liu, 2004) is one of the most popular packages written in the Maple programming language, which is a comprehensive and complete implementation of some powerful methods such as the tanh-method, the extended tanh-method, the Jacobi elliptic function method, and the elliptic equation method. One can note that most packages have been developed within commercially available software frameworks, such as Maple and Mathematica. Besides this, all these computer packages have implemented the function-expansion methods. One serious drawback of the function-expansion method is that the solutions which contain functions other than some specific type of functions, such as tanh, sech, JacobiSN, JacobiCN, are not obtained. Additionally, the non-polynomial forms of these particular functions are not obtained also. On the other hand, we have observed that F-expansion, mF-expansion, and FIM are different kinds of methods, which can overcome the limitations of the function-expansion method. To the best of our knowledge, the computer packages implementing F-expansion and first integral methods are not available so far. Keeping in mind all the above points of view, we have been motivated to develop a free and open-source computer package or C++ library called GiNaCDE, which implements the F-expansion and first integral methods.

The symbolic manipulations of GiNaCDE depend only on GiNaC (Bauer et al., 2002). There are several advantages to use GiNaC over other CAS. GiNaC is a free and open-source pure C++ library. It can accept C++ programming language, a general-purpose object-oriented programming (OOP) language, and it is fast like commercially available computer algebra systems. Besides the library version of GiNaCDE, we have also developed a GUI version of GiNaCDE called GiNaCDE GUI, which facilitates users to solve NLPDEs automatically without writing programming and compilation each time. This GUI version guides us in each step to obtain the output results.

References

- Baldwin, D., Göktaş, Ü., Hereman, W., Hong, L., Martino, R. S., & Miller, J. C. (2004). Symbolic computation of exact solutions expressible in hyperbolic and elliptic functions for nonlinear PDEs. *Journal of Symbolic Computation*, 37(6), 669–705. <https://doi.org/10.1016/j.jsc.2003.09.004>
- Bauer, C., Frink, A., & Kreckel, R. (2002). Introduction to the GiNaC framework for symbolic computation within the c++ programming language. *Journal of Symbolic Computation*, 33(1), 1–12. <https://doi.org/10.1006/jasco.2001.0494>

¹<https://github.com/mithun218/GiNaCDE/blob/master/doc/documentation.pdf>

- Bhrawy, A., Abdelkawy, M., Kumar, Dr. S., & Biswas, A. (2013). Solitons and other solutions to kadomtsev-petviashvili equation of b-type. *Romanian Journal of Physics*, 58(7-8), 729–748.
- Cai, G., Wang, Q., & Huang, J. (2006). A modified f-expansion method for solving breaking soliton equation. *International Journal of Nonlinear Science*, 2, 122–128.
- Feng, Z. (2002). On explicit exact solutions to the compound burgers-KdV equation. *Physics Letters A*, 293, 57–66. [https://doi.org/10.1016/S0375-9601\(01\)00825-8](https://doi.org/10.1016/S0375-9601(01)00825-8)
- He, Y. (2013). New jacobi elliptic function solutions for the kudryashov-sinelshchikov equation using improved f-expansion method. *Mathematical Problems in Engineering*, 2013. <https://doi.org/10.1155/2013/104894>
- Huang, D., Li, D., & Zhang, H. (2007). Explicit and exact traveling solutions for the generalized derivative schrödinger equation. *Chaos, Solitons & Fractals*, 31, 586–593. <https://doi.org/10.1016/j.chaos.2005.10.007>
- Li, Z., & Liu, Y. (2002). RATH: A maple package for finding travelling solitary wave solutions to nonlinear evolution equations. *Computer Physics Communications*, 148(2), 256–266. [https://doi.org/10.1016/S0010-4655\(02\)00559-3](https://doi.org/10.1016/S0010-4655(02)00559-3)
- Lin, Z.-B., & Liu, Y.-P. (2004). RAEEM: A maple package for finding a series of exact traveling wave solutions for nonlinear evolution equations. *Computer Physics Communications*, 163, 191–201. <https://doi.org/10.1016/j.cpc.2004.08.007>
- Mirzazadeh, M., & Eslami, M. (2012). Exact solutions of the kudryashov-sinelshchikov equation and nonlinear telegraph equation via the first integral method. *Nonlinear Analysis. Modelling and Control*, 4. <https://doi.org/10.15388/NA.17.4.14052>
- Na, S. (2004). New exact travelling wave solutions for the kawahara and modified kawahara equations. *Chaos, Solitons & Fractals*, 19, 147–150. [https://doi.org/10.1016/S0960-0779\(03\)00102-4](https://doi.org/10.1016/S0960-0779(03)00102-4)
- Parkes, E. J., & Duffy, B. R. (1996). An automated tanh-function method for finding solitary wave solutions to non-linear evolution equations. *Computer Physics Communications*, 98(3), 288–300. [https://doi.org/10.1016/0010-4655\(96\)00104-X](https://doi.org/10.1016/0010-4655(96)00104-X)
- Zhang, Z., Liu, Z., Miao, X., & Chen, Y. (2010). New exact solutions to the perturbed nonlinear schrödinger's equation with kerr law nonlinearity. *Applied Mathematics and Computation*, 216(10), 3064–3072. <https://doi.org/10.1016/j.amc.2010.04.026>
- Zhou, Y., Wang, M., & Wang, Y. (2003). Periodic wave solutions to a coupled KdV equations with variable coefficients. *Physics Letters A*, 308(1), 31–36. [https://doi.org/10.1016/S0375-9601\(02\)01775-9](https://doi.org/10.1016/S0375-9601(02)01775-9)